


Fourth Quarter 2009

Vol. 7, No. 4



HIV/AIDS Surveillance Program

M. Beth Scalco, LCSW MPA

HIV/AIDS Program Administrative Director

Amy Zapata, MPH

HIV/AIDS Surveillance Program Manager

Editor/Production:

Jessica Fridge

William T. Robinson

Data Management and Analysis:

Debbie Wendell, MPH, Ph.D., *Coordinator*

Jessica Fridge, MSPH, *Surveillance Data Supervisor*

Tania Bechtel, *Data Manager*

Joseph Foxhood, *Information Systems Developer*

George Stephens, *Computer Technician*

Josette Russell, *Data Manager*

Special Projects:

William T. Robinson, Ph.D., *Biostatistician*

Jessica Cole, MPH, *Perinatal Surveillance Coordinator*

Field Epidemiologists:

Betsy Shepard, PAC, MPH, *Epidemiologist Supervisor*

James Hubbard **Region 1**
Connie Gordon **Regions 1, 3**
Tanya Brown, RHIA **Region 2**

Cheryl Edney, BS MT(ASCP) **Regions 4, 5, 6**
Gayle Norris, MA **Regions 7, 8**
Gale Terry, LPN **Regions 2, 9**

TABLE OF CONTENTS

| Contents | Page |
|---|-------------|
| Surveillance Summary | 1 |
| Louisiana: Persons Living with HIV/AIDS | 2 |
| Persons Living with AIDS | 3 |
| Persons newly diagnosed with HIV/AIDS | 4 |
| Parish: Persons Living with HIV/AIDS | 5 |
| Persons Living with AIDS | 6 |
| Metro New Orleans: Persons Living with HIV/AIDS | 7 |
| Persons Living with AIDS | 8 |
| Persons newly diagnosed with HIV/AIDS | 9 |
| Region 1: Persons Living with HIV/AIDS | 10 |
| Persons Living with AIDS | 11 |
| Persons newly diagnosed with HIV/AIDS | 12 |
| Region 2: Persons Living with HIV/AIDS | 13 |
| Persons Living with AIDS | 14 |
| Persons newly diagnosed with HIV/AIDS | 15 |
| Region 3: Persons Living with HIV/AIDS | 16 |
| Persons Living with AIDS | 17 |
| Persons newly diagnosed with HIV/AIDS | 18 |
| Region 4: Persons Living with HIV/AIDS | 19 |
| Persons Living with AIDS | 20 |
| Persons newly diagnosed with HIV/AIDS | 21 |
| Region 5: Persons Living with HIV/AIDS | 22 |
| Persons Living with AIDS | 23 |
| Persons newly diagnosed with HIV/AIDS | 24 |
| Region 6: Persons Living with HIV/AIDS | 25 |
| Persons Living with AIDS | 26 |
| Persons newly diagnosed with HIV/AIDS | 27 |
| Region 7: Persons Living with HIV/AIDS | 28 |
| Persons Living with AIDS | 29 |
| Persons newly diagnosed with HIV/AIDS | 30 |
| Region 8: Persons Living with HIV/AIDS | 31 |
| Persons Living with AIDS | 32 |
| Persons newly diagnosed with HIV/AIDS | 33 |
| Region 9: Persons Living with HIV/AIDS | 34 |
| Persons Living with AIDS | 35 |
| Persons newly diagnosed with HIV/AIDS | 36 |

Surveillance Summary

Data in this report includes cases diagnosed with HIV/AIDS and reported as of December 31, 2009¹.

- As of December 31, 2009, a cumulative total of **29,866** HIV/AIDS cases have been reported in Louisiana, including **312** pediatric cases.
- The cumulative number of deaths among persons diagnosed with HIV/AIDS, reported to the Louisiana Office of Public Health through December 31, 2009, is **12,530** including **101** deaths among pediatric cases.
- A total of **17,387** persons are living with HIV/AIDS in Louisiana; of these individuals, **9,379** persons (54%) have an AIDS diagnosis.
- Among women living with HIV/AIDS, the predominant known mode of transmission is high-risk heterosexual contact followed by injection drug use. Among men living with HIV/AIDS, the predominant mode of transmission is men who have sex with men followed by high-risk heterosexual contact.
- Nationally, Louisiana ranked **5th** highest in AIDS case rates and **11th** in the number of AIDS cases diagnosed in 2007, according to the CDC 2007 HIV/AIDS Surveillance Report (Vol. 19).
- New Orleans ranked **2nd** for AIDS case rates among the largest metropolitan areas in the U.S. in 2007; Baton Rouge ranked **3rd**.
- In 2008, **1,158** new HIV cases were diagnosed in Louisiana. Of these cases, **33%** (n=385) occurred in Region 1 (New Orleans) and **26%** (n=301) occurred in Region 2 (Baton Rouge).
- In 2008, **865** new AIDS cases were diagnosed in Louisiana. Of these cases, **31%** (n=272) occurred in Region 1 (New Orleans) and **29%** (n=251) occurred in Region 2 (Baton Rouge).
- In Louisiana, **33%** of new HIV cases and **34%** of new AIDS cases are among women.
- HIV continues to disproportionately affect African Americans in Louisiana. In 2008, **72%** of newly diagnosed HIV cases and **71%** of newly diagnosed AIDS cases were among African Americans.

¹Please note: This report reflects only information reported to Louisiana's HIV/AIDS surveillance system and information on persons living with HIV/AIDS (PLWH/A) may not be accurate due to population changes resulting from Hurricane Katrina. In March 2006, the Louisiana Office of Public Health HIV/AIDS Program (OPH-HAP) published the Louisiana HIV/AIDS Surveillance First Quarter 2006 report. In addition to the HIV/AIDS surveillance information, as presented here, this report contained a description of two methodologies used to estimate the impact of Hurricane Katrina on the number of PLWH/A in the New Orleans metropolitan area (see <http://www.dhh.louisiana.gov/offices/publications/pubs-264/1st%20Quarter%202006.pdf> for the full report). Updated estimates were provided in the 2006 Third Quarter Report at: <http://www.dhh.louisiana.gov/offices/publications/pubs-264/Third%20Quarter%202006.pdf>. Additional updated population estimates were presented at the February 2007 Statewide HIV/AIDS Meeting. This presentation can be found at: <http://www.dhh.louisiana.gov/offices/page.asp?ID=264&Detail=7584>. Updated estimates were also presented in the 2005-2006 Annual Report located at <http://www.dhh.louisiana.gov/offices/reports.asp?ID=264&Detail=261>. More recent estimates indicate that as much as 84% of the pre-Katrina PLWH/A may now be living within the metropolitan area.

Persons Living with HIV/AIDS Louisiana

- **29,866** cumulative cases have been detected in Louisiana, including 312 pediatric cases.
- **12,530** cumulative deaths among persons with HIV/AIDS have occurred in Louisiana, including 101 pediatric deaths.
- **17,387** persons are currently living with HIV/AIDS in Louisiana.
 - Persons living with HIV: 8,008
 - Persons living with AIDS: 9,379

Age Category

| Current Age | Cases | Percent |
|--------------------|---------------|----------------|
| 0 - 12 | 78 | <1 % |
| 13 - 19 | 187 | 1 % |
| 20 - 24 | 772 | 4 % |
| 25 - 29 | 1,453 | 8 % |
| 30 - 34 | 1,925 | 11 % |
| 35 - 39 | 2,178 | 13 % |
| 40 - 44 | 2,762 | 16 % |
| 45 - 49 | 3,177 | 18 % |
| 50 - 54 | 2,329 | 13 % |
| 55 - 59 | 1,356 | 8 % |
| 60+ | 1,170 | 7 % |
| Total | 17,387 | 100 % |

Race/Ethnicity

| | Men | Women | Total | Percent |
|------------------------|---------------|--------------|---------------|----------------|
| Black, Non-Hispanic | 7,318 | 4,262 | 11,580 | 67 % |
| White, Non-Hispanic | 4,240 | 763 | 5,003 | 29 % |
| Hispanic | 517 | 112 | 629 | 4 % |
| Asian/Pacific Islander | 45 | 11 | 56 | <1 % |
| American Indian | 19 | 6 | 25 | <1 % |
| Unknown | 17 | 4 | 21 | <1 % |
| Multi-race | 52 | 21 | 73 | <1 % |
| Total | 12,208 | 5,179 | 17,387 | 100 % |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|--------------|--------------|---------------|----------------|
| Men who have sex with men | 5,324 | 0 | 5,324 | 48 % |
| Injection drug use | 1,167 | 686 | 1,853 | 17 % |
| Men who have sex with men and inject drugs | 874 | 0 | 874 | 8 % |
| Heterosexual contact | 890 | 1,869 | 2,759 | 25 % |
| Transfusion/Hemophilia | 65 | 39 | 104 | 1 % |
| Confirmed Other | 0 | 1 | 1 | <1 % |
| Perinatal | 95 | 93 | 188 | 2 % |
| Total | 8,415 | 2,688 | 11,103 | 100 % |
| <i>No Identified Risk</i> | <i>3,793</i> | <i>2,491</i> | <i>6,284</i> | |

Persons Living with AIDS Louisiana

- **20,143** cumulative AIDS cases have been diagnosed in Louisiana, including 144 pediatric cases.
- **9,379** persons are currently living with AIDS in Louisiana.

Age Category

| Current Age | Cases | Percent |
|--------------------|--------------|----------------|
| 0 - 12 | 9 | <1 % |
| 13 - 19 | 51 | 1 % |
| 20 - 24 | 187 | 2 % |
| 25 - 29 | 511 | 5 % |
| 30 - 34 | 824 | 9 % |
| 35 - 39 | 1,131 | 12 % |
| 40 - 44 | 1,644 | 18 % |
| 45 - 49 | 1,951 | 21 % |
| 50 - 54 | 1,455 | 16 % |
| 55 - 59 | 871 | 9 % |
| 60+ | 745 | 8 % |
| Total | 9,379 | 100 % |

Race/Ethnicity

| | Men | Women | Total | Percent |
|---------------------|--------------|--------------|--------------|----------------|
| Black, Non-Hispanic | 4,053 | 2,063 | 6,116 | 65 % |
| White, Non- | 2,485 | 349 | 2,834 | 30 % |
| Hispanic | 279 | 68 | 347 | 4 % |
| Asian/Pacific | 18 | 5 | 23 | <1 % |
| American Indian | 9 | 3 | 12 | <1 % |
| Unknown | 2 | 0 | 2 | <1 % |
| Multi-race | 34 | 11 | 45 | <1 % |
| Total | 6,880 | 2,499 | 9,379 | 100 % |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|--------------|--------------|--------------|----------------|
| Men who have sex with men | 3,077 | 0 | 3,077 | 47 % |
| Injection drug use | 775 | 403 | 1,178 | 18 % |
| Men who have sex with men and inject drugs | 594 | 0 | 594 | 9 % |
| Heterosexual contact | 519 | 1,002 | 1,521 | 23 % |
| Transfusion/Hemophilia | 46 | 24 | 70 | 1 % |
| Perinatal | 25 | 30 | 55 | 1 % |
| Total | 5036 | 1459 | 6,495 | 100 % |
| <i>No Identified Risk</i> | <i>1,844</i> | <i>1,040</i> | <i>2,884</i> | |

**Persons Newly Diagnosed with HIV/AIDS
Louisiana**

HIV CASES DIAGNOSED January - December 2008

• 1,158 persons were newly diagnosed with HIV in Louisiana.

| Gender | Cases | Percent | Race/Ethnicity | Cases | Percent |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 772 | 67 % | Black, Non-Hispanic | 838 | 72 % |
| Women | 386 | 33 % | White, Non-Hispanic | 231 | 20 % |
| Total | 1158 | 100 % | Hispanic | 66 | 6 % |
| | | | Asian/Pacific Islander | 5 | <1 % |
| | | | American Indian | 1 | <1 % |
| | | | Unknown | 4 | <1 % |
| | | | Multi-race | 13 | 1 % |
| | | | Total | 1,158 | 100 % |

| Age at Detection | Cases | Percent | Cases with a reported risk: | Cases | Percent |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 0 - 12 | 5 | <1 % | Men who have sex with men | 338 | 60 % |
| 13 - 24 | 247 | 21 % | Injection drug use | 49 | 9 % |
| 25 - 34 | 344 | 30 % | Men who have sex with men and inject drugs | 10 | 2 % |
| 35 - 44 | 249 | 22 % | Heterosexual contact | 160 | 28 % |
| 45+ | 313 | 27 % | Perinatal | 5 | 1 % |
| Total | 1,158 | 100 % | Total | 562 | 100 % |
| | | | <i>No Identified Risk</i> | <i>596</i> | |

AIDS CASES DIAGNOSED January - December 2008

• 865 persons were newly diagnosed with AIDS in Louisiana.

| Gender | Cases | Percent | Race/Ethnicity | Cases | Percent |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 571 | 66 % | Black, Non-Hispanic | 610 | 71 % |
| Women | 294 | 34 % | White, Non-Hispanic | 207 | 24 % |
| Total | 865 | 100 % | Hispanic | 40 | 5 % |
| | | | Asian/Pacific Islander | 3 | <1 % |
| | | | Multi-race | 5 | 1 % |
| | | | Total | 865 | 100 % |

| Age at Diagnosis | Cases | Percent | Cases with a reported risk: | Cases | Percent |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 0 - 12 | 1 | <1 % | Men who have sex with men | 221 | 47 % |
| 13 - 24 | 66 | 8 % | Injection drug use | 84 | 18 % |
| 25 - 34 | 235 | 27 % | Men who have sex with men and inject drugs | 20 | 4 % |
| 35 - 44 | 254 | 29 % | Heterosexual contact | 140 | 30 % |
| 45+ | 309 | 36 % | Transfusion/Hemophilia | 3 | 1 % |
| Total | 865 | 100 % | Perinatal | 4 | 1 % |
| | | | Total | 472 | 100 % |
| | | | <i>No Identified Risk</i> | <i>393</i> | |

Persons Living with HIV/AIDS by Parish

Fourth Quarter 2009

| <u>Parish</u> | <u>Living Cases</u> | <u>Parish</u> | <u>Living Cases</u> |
|------------------|---------------------|----------------------|---------------------|
| Acadia | 89 | Sabine | 22 |
| Allen | 237 | St. Bernard | 105 |
| Ascension | 172 | St. Charles | 82 |
| Assumption | 23 | St. Helena | 19 |
| Avoyelles | 172 | St. James | 63 |
| Beauregard | 35 | St. John the Baptist | 105 |
| Bienville | 30 | St. Landry | 242 |
| Bossier | 171 | St. Martin | 100 |
| Caddo | 919 | St. Mary | 79 |
| Calcasieu | 598 | St. Tammany | 334 |
| Caldwell | 48 | Tangipahoa | 282 |
| Cameron | 5 | Tensas | 35 |
| Catahoula | 24 | Terrebonne | 183 |
| Claiborne | 78 | Union | 41 |
| Concordia | 39 | Vermilion | 79 |
| De Soto | 55 | Vernon | 44 |
| East Baton Rouge | 3,263 | Washington | 181 |
| East Carroll | 40 | Webster | 49 |
| East Feliciana | 139 | West Baton Rouge | 109 |
| Evangeline | 57 | West Carroll | 10 |
| Franklin | 26 | West Feliciana | 145 |
| Grant | 28 | Winn | 94 |
| Iberia | 104 | | |
| Iberville | 301 | | |
| Jackson | 36 | | |
| Jefferson | 1,605 | | |
| Jefferson Davis | 53 | | |
| La Salle | 19 | | |
| Lafayette | 607 | | |
| Lafourche | 100 | | |
| Lincoln | 61 | | |
| Livingston | 146 | | |
| Madison | 59 | | |
| Morehouse | 59 | | |
| Natchitoches | 95 | | |
| Orleans | 4,465 | | |
| Ouachita | 496 | | |
| Plaquemines | 31 | | |
| Pointe Coupee | 47 | | |
| Rapides | 402 | | |
| Red River | 8 | | |
| Richland | 39 | | |
| | | Total | 17,387 |

Persons Living with AIDS by Parish

Fourth Quarter 2009

| <u>Parish</u> | <u>Living Cases</u> | <u>Parish</u> | <u>Living Cases</u> |
|------------------|---------------------|----------------------|---------------------|
| Acadia | 50 | Sabine | 16 |
| Allen | 115 | St. Bernard | 62 |
| Ascension | 89 | St. Charles | 46 |
| Assumption | 16 | St. Helena | 10 |
| Avoyelles | 94 | St. James | 32 |
| Beauregard | 22 | St. John the Baptist | 61 |
| Bienville | 14 | St. Landry | 143 |
| Bossier | 96 | St. Martin | 57 |
| Caddo | 511 | St. Mary | 51 |
| Calcasieu | 352 | St. Tammany | 191 |
| Caldwell | 22 | Tangipahoa | 156 |
| Cameron | 3 | Tensas | 17 |
| Catahoula | 14 | Terrebonne | 103 |
| Claiborne | 45 | Union | 27 |
| Concordia | 22 | Vermilion | 46 |
| De Soto | 33 | Vernon | 21 |
| East Baton Rouge | 1,687 | Washington | 100 |
| East Carroll | 25 | Webster | 22 |
| East Feliciana | 80 | West Baton Rouge | 63 |
| Evangeline | 30 | West Carroll | 7 |
| Franklin | 12 | West Feliciana | 93 |
| Grant | 15 | Winn | 49 |
| Iberia | 57 | | |
| Iberville | 167 | | |
| Jackson | 21 | | |
| Jefferson | 873 | | |
| Jefferson Davis | 30 | | |
| La Salle | 11 | | |
| Lafayette | 310 | | |
| Lafourche | 63 | | |
| Lincoln | 30 | | |
| Livingston | 81 | | |
| Madison | 29 | | |
| Morehouse | 28 | | |
| Natchitoches | 51 | | |
| Orleans | 2,363 | | |
| Ouachita | 266 | | |
| Plaquemines | 22 | | |
| Pointe Coupee | 30 | | |
| Rapides | 194 | | |
| Red River | 7 | | |
| Richland | 24 | | |

Total **9,379**

Persons Living with HIV/AIDS Metro New Orleans

Jefferson, Orleans, St. Tammany, St. John the Baptist, Plaquemines, St. Bernard, St. James, and St. Charles Parishes

- **14,259** cumulative cases have been detected in the New Orleans MSA, including 152 pediatric cases.
- **6,045** cumulative deaths among persons with HIV/AIDS have occurred in the New Orleans MSA, including 51 pediatric deaths
- **6,790** persons are currently living with HIV/AIDS in the New Orleans MSA.
Persons living with HIV: 3,140
Persons living with AIDS: 3,650

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|--------------|--------------|------------------------|--------------|--------------|--------------|--------------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 29 | <1 % | Black, Non-Hispanic | 2,534 | 1,444 | 3,978 | 59 % |
| 13 - 19 | 60 | 1 % | White, Non-Hispanic | 2,142 | 243 | 2,385 | 35 % |
| 20 - 24 | 270 | 4 % | Hispanic | 266 | 80 | 346 | 5 % |
| 25 - 29 | 474 | 7 % | Asian/Pacific Islander | 29 | 7 | 36 | 1 % |
| 30 - 34 | 661 | 10 % | American Indian | 6 | 1 | 7 | <1 % |
| 35 - 39 | 819 | 12 % | Unknown | 10 | 1 | 11 | <1 % |
| 40 - 44 | 1,073 | 16 % | Multi-race | 21 | 6 | 27 | <1 % |
| 45 - 49 | 1,284 | 19 % | Total | 5,008 | 1,782 | 6,790 | 100 % |
| 50 - 54 | 959 | 14 % | | | | | |
| 55 - 59 | 604 | 9 % | | | | | |
| 60+ | 557 | 8 % | | | | | |
| Total | 6,790 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|--------------|------------|--------------|--------------|
| Men who have sex with men | 2,497 | 0 | 2,497 | 57 % |
| Injection drug use | 355 | 238 | 593 | 13 % |
| Men who have sex with men and inject drugs | 332 | 0 | 332 | 8 % |
| Heterosexual contact | 254 | 612 | 866 | 20 % |
| Transfusion/Hemophilia | 27 | 19 | 46 | 1 % |
| Perinatal | 30 | 37 | 67 | 2 % |
| Total | 3,495 | 906 | 4,401 | 100 % |
| <i>No Identified Risk</i> | <i>1,513</i> | <i>876</i> | <i>2,389</i> | |

Persons Living with AIDS

Metro New Orleans

Jefferson, Orleans, St. Tammany, St. John the Baptist, Plaquemines, St. Bernard, St. James, and St. Charles Parishes

- **9,747** cumulative AIDS cases have been diagnosed in the New Orleans MSA, including 71 pediatric cases.
- **3,650** persons are currently living with AIDS in the New Orleans MSA.

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|-------|---------|------------------------|-------|-------|-------|---------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 4 | <1 % | Black, Non-Hispanic | 1,350 | 719 | 2,069 | 57 % |
| 13 - 19 | 16 | <1 % | White, Non-Hispanic | 1,232 | 99 | 1,331 | 36 % |
| 20 - 24 | 55 | 2 % | Hispanic | 159 | 53 | 212 | 6 % |
| 25 - 29 | 161 | 4 % | Asian/Pacific Islander | 11 | 3 | 14 | <1 % |
| 30 - 34 | 261 | 7 % | American Indian | 3 | 1 | 4 | <1 % |
| 35 - 39 | 415 | 11 % | Unknown | 2 | 0 | 2 | <1 % |
| 40 - 44 | 613 | 17 % | Multi-race | 14 | 4 | 18 | <1 % |
| 45 - 49 | 796 | 22 % | Total | 2,771 | 879 | 3,650 | 100 % |
| 50 - 54 | 595 | 16 % | | | | | |
| 55 - 59 | 375 | 10 % | | | | | |
| 60+ | 359 | 10 % | | | | | |
| Total | 3,650 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|-------|-------|-------|---------|
| Men who have sex with men | 1,431 | 0 | 1,431 | 56 % |
| Injection drug use | 230 | 134 | 364 | 14 % |
| Men who have sex with men and inject drugs | 222 | 0 | 222 | 9 % |
| Heterosexual contact | 159 | 342 | 501 | 20 % |
| Transfusion/Hemophilia | 19 | 10 | 29 | 1 % |
| Perinatal | 8 | 13 | 21 | 1 % |
| Total | 2,069 | 499 | 2,568 | 100 % |
| <i>No Identified Risk</i> | 702 | 380 | 1,082 | |

Persons Newly Diagnosed with HIV/AIDS

Metro New Orleans

Jefferson, Orleans, St. Tammany, St. John the Baptist, Plaquemines, St. Bernard, St. James, and St. Charles Parishes

HIV CASES DIAGNOSED January - December 2008

- 422 persons were newly diagnosed with HIV in the New Orleans MSA.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 294 | 70 % | Black, Non-Hispanic | 272 | 64 % |
| Women | 128 | 30 % | White, Non-Hispanic | 101 | 24 % |
| Total | 422 | 100 % | Hispanic | 37 | 9 % |
| | | | Asian/Pacific Islander | 3 | 1 % |
| | | | Unknown | 2 | <1 % |
| | | | Multi-race | 7 | 2 % |
| | | | Total | 422 | 100 % |

| <u>Age at Detection</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 0 - 12 | 3 | 1 % | Men who have sex with men | 149 | 66 % |
| 13 - 24 | 89 | 21 % | Injection drug use | 20 | 9 % |
| 25 - 34 | 130 | 31 % | Men who have sex with men and inject drugs | 5 | 2 % |
| 35 - 44 | 96 | 23 % | Heterosexual contact | 49 | 22 % |
| 45+ | 104 | 25 % | Perinatal | 3 | 1 % |
| Total | 422 | 100 % | Total | 226 | 100 % |
| | | | <i>No Identified Risk</i> | <i>196</i> | |

AIDS CASES DIAGNOSED January - December 2008

- 290 persons were newly diagnosed with AIDS in the New Orleans MSA.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 200 | 69 % | Black, Non-Hispanic | 183 | 63 % |
| Women | 90 | 31 % | White, Non-Hispanic | 79 | 27 % |
| Total | 290 | 100 % | Hispanic | 27 | 9 % |
| | | | Multi-race | 1 | <1 % |
| | | | Total | 290 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 0 - 12 | 1 | <1 % | Men who have sex with men | 83 | 52 % |
| 13 - 24 | 19 | 7 % | Injection drug use | 23 | 14 % |
| 25 - 34 | 79 | 27 % | Men who have sex with men and inject drugs | 7 | 4 % |
| 35 - 44 | 86 | 30 % | Heterosexual contact | 46 | 29 % |
| 45+ | 105 | 36 % | Transfusion/Hemophilia | 1 | 1 % |
| Total | 290 | 100 % | Perinatal | 1 | 1 % |
| | | | Total | 161 | 100 % |
| | | | <i>No Identified Risk</i> | <i>129</i> | |

**Persons Living with HIV/AIDS
Region 1: New Orleans Region**

- **13,434** cumulative cases have been detected in Region 1, including **136** pediatric cases.
- **5,690** cumulative deaths among persons in Region 1 with HIV/AIDS have occurred, including **46** pediatric deaths.
- **6,206** persons are currently living with HIV/AIDS in Region 1.
Persons living with HIV: **2,886**
Persons living with AIDS: **3,320**

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|--------------|--------------|------------------------|--------------|--------------|--------------|--------------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 21 | <1 % | Black, Non-Hispanic | 2,366 | 1,337 | 3,703 | 60 % |
| 13 - 19 | 55 | 1 % | White, Non-Hispanic | 1,903 | 192 | 2,095 | 34 % |
| 20 - 24 | 242 | 4 % | Hispanic | 254 | 77 | 331 | 5 % |
| 25 - 29 | 431 | 7 % | Asian/Pacific Islander | 28 | 7 | 35 | 1 % |
| 30 - 34 | 612 | 10 % | American Indian | 6 | 1 | 7 | <1 % |
| 35 - 39 | 735 | 12 % | Unknown | 10 | 1 | 11 | <1 % |
| 40 - 44 | 981 | 16 % | Multi-race | 18 | 6 | 24 | <1 % |
| 45 - 49 | 1,168 | 19 % | Total | 4,585 | 1,621 | 6,206 | 100 % |
| 50 - 54 | 882 | 14 % | | | | | |
| 55 - 59 | 565 | 9 % | | | | | |
| 60+ | 514 | 8 % | | | | | |
| Total | 6,206 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|--------------|------------|--------------|--------------|
| Men who have sex with men | 2,277 | 0 | 2,277 | 57 % |
| Injection drug use | 325 | 217 | 542 | 14 % |
| Men who have sex with men and inject drugs | 316 | 0 | 316 | 8 % |
| Heterosexual contact | 226 | 557 | 783 | 20 % |
| Transfusion/Hemophilia | 24 | 18 | 42 | 1 % |
| Perinatal | 24 | 29 | 53 | 1 % |
| Total | 3,192 | 821 | 4,013 | 100 % |
| <i>No Identified Risk</i> | <i>1,393</i> | <i>800</i> | <i>2,193</i> | |

Persons Living with AIDS
Region 1: New Orleans Region

- **9,157** cumulative AIDS cases have been diagnosed in Region 1, including **63** pediatric cases.
- **3,320** persons are currently living with AIDS in Region 1.

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|-------|---------|------------------------|-------|-------|-------|---------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 4 | <1 % | Black, Non-Hispanic | 1,245 | 666 | 1,911 | 58 % |
| 13 - 19 | 13 | <1 % | White, Non-Hispanic | 1,092 | 77 | 1,169 | 35 % |
| 20 - 24 | 51 | 2 % | Hispanic | 152 | 52 | 204 | 6 % |
| 25 - 29 | 149 | 4 % | Asian/Pacific Islander | 11 | 3 | 14 | <1 % |
| 30 - 34 | 238 | 7 % | American Indian | 3 | 1 | 4 | <1 % |
| 35 - 39 | 369 | 11 % | Unknown | 2 | 0 | 2 | <1 % |
| 40 - 44 | 555 | 17 % | Multi-race | 12 | 4 | 16 | <1 % |
| 45 - 49 | 719 | 22 % | Total | 2,517 | 803 | 3,320 | 100 % |
| 50 - 54 | 546 | 16 % | | | | | |
| 55 - 59 | 348 | 10 % | | | | | |
| 60+ | 328 | 10 % | | | | | |
| Total | 3,320 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|------|-------|-------|---------|
| Men who have sex with men | 1304 | 0 | 1304 | 56 % |
| Injection drug use | 209 | 122 | 331 | 14 % |
| Men who have sex with men and inject drugs | 209 | 0 | 209 | 9 % |
| Heterosexual contact | 139 | 312 | 451 | 19 % |
| Transfusion/Hemophilia | 16 | 9 | 25 | 1 % |
| Perinatal | 6 | 11 | 17 | 1 % |
| Total | 1883 | 454 | 2337 | 100 % |
| <i>No Identified Risk</i> | 634 | 349 | 983 | |

Persons Newly Diagnosed with HIV/AIDS

Region 1: New Orleans Region

HIV CASES DIAGNOSED January - December 2008

- 385 persons were newly diagnosed with HIV in Region 1.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 271 | 70 % | Black, Non-Hispanic | 252 | 65 % |
| Women | 114 | 30 % | White, Non-Hispanic | 86 | 22 % |
| Total | 385 | 100 % | Hispanic | 35 | 9 % |
| | | | Asian/Pacific Islander | 3 | 1 % |
| | | | Unknown | 2 | 1 % |
| | | | Multi-race | 7 | 2 % |
| | | | Total | 385 | 100 % |

| <u>Age at Detection</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 0 - 12 | 2 | 1 % | Men who have sex with men | 139 | 66 % |
| 13 - 24 | 83 | 22 % | Injection drug use | 18 | 9 % |
| 25 - 34 | 116 | 30 % | Men who have sex with men and inject drugs | 4 | 2 % |
| 35 - 44 | 84 | 22 % | Heterosexual contact | 48 | 23 % |
| 45+ | 100 | 26 % | Perinatal | 2 | 1 % |
| Total | 385 | 100 % | Total | 211 | 100 % |

| | |
|---------------------------|------------|
| <i>No Identified Risk</i> | <i>174</i> |
|---------------------------|------------|

AIDS CASES DIAGNOSED January - December 2008

- 272 persons were newly diagnosed with AIDS in Region 1.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 187 | 69 % | Black, Non-Hispanic | 172 | 63 % |
| Women | 85 | 31 % | White, Non-Hispanic | 73 | 27 % |
| Total | 272 | 100 % | Hispanic | 26 | 10 % |
| | | | Multi-race | 1 | <1 % |
| | | | Total | 272 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 0 - 12 | 1 | <1 % | Men who have sex with men | 79 | 52 % |
| 13 - 24 | 18 | 7 % | Injection drug use | 21 | 14 % |
| 25 - 34 | 73 | 27 % | Men who have sex with men and inject drugs | 6 | 4 % |
| 35 - 44 | 79 | 29 % | Heterosexual contact | 45 | 29 % |
| 45+ | 101 | 37 % | Transfusion/Hemophilia | 1 | 1 % |
| Total | 272 | 100 % | Perinatal | 1 | 1 % |
| | | | Total | 153 | 100 % |

| | |
|---------------------------|------------|
| <i>No Identified Risk</i> | <i>119</i> |
|---------------------------|------------|

Persons Living with HIV/AIDS
Region 2: Baton Rouge Region

- **6,491** cumulative cases have been detected in Region 2, including **64** pediatric cases.
- **2,712** cumulative deaths among persons in Region 2 with HIV/AIDS have occurred, including **15** pediatric deaths.
- **4,176** persons are currently living with HIV/AIDS in Region 2.
Persons living with HIV: **1,967**
Persons living with AIDS: **2,209**

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|--------------|--------------|------------------------|--------------|--------------|--------------|--------------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 25 | 1 % | Black, Non-Hispanic | 2,021 | 1,456 | 3,477 | 83 % |
| 13 - 19 | 52 | 1 % | White, Non-Hispanic | 514 | 119 | 633 | 15 % |
| 20 - 24 | 225 | 5 % | Hispanic | 36 | 11 | 47 | 1 % |
| 25 - 29 | 412 | 10 % | Asian/Pacific Islander | 5 | 1 | 6 | <1 % |
| 30 - 34 | 526 | 13 % | American Indian | 2 | 0 | 2 | <1 % |
| 35 - 39 | 546 | 13 % | Unknown | 2 | 0 | 2 | <1 % |
| 40 - 44 | 601 | 14 % | Multi-race | 7 | 2 | 9 | <1 % |
| 45 - 49 | 714 | 17 % | Total | 2,587 | 1,589 | 4,176 | 100 % |
| 50 - 54 | 514 | 12 % | | | | | |
| 55 - 59 | 308 | 7 % | | | | | |
| 60+ | 253 | 6 % | | | | | |
| Total | 4,176 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|--------------|------------|--------------|--------------|
| Men who have sex with men | 856 | 0 | 856 | 35 % |
| Injection drug use | 371 | 226 | 597 | 24 % |
| Men who have sex with men and inject drugs | 169 | 0 | 169 | 7 % |
| Heterosexual contact | 242 | 527 | 769 | 31 % |
| Transfusion/Hemophilia | 8 | 11 | 19 | 1 % |
| Perinatal | 28 | 22 | 50 | 2 % |
| Total | 1,674 | 786 | 2,460 | 100 % |
| <i>No Identified Risk</i> | <i>913</i> | <i>803</i> | <i>1,716</i> | |

Persons Living with AIDS
Region 2: Baton Rouge Region

- **4,323** cumulative AIDS cases have been diagnosed in Region 2, including **22** pediatric cases.
- **2,209** persons are currently living with AIDS in Region 2.

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|-------|---------|------------------------|-------|-------|-------|---------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 13 - 19 | 17 | 1 % | Black, Non-Hispanic | 1,122 | 707 | 1,829 | 83 % |
| 20 - 24 | 61 | 3 % | White, Non-Hispanic | 292 | 51 | 343 | 16 % |
| 25 - 29 | 146 | 7 % | Hispanic | 20 | 6 | 26 | 1 % |
| 30 - 34 | 226 | 10 % | Asian/Pacific Islander | 2 | 0 | 2 | <1 % |
| 35 - 39 | 285 | 13 % | American Indian | 2 | 0 | 2 | <1 % |
| 40 - 44 | 362 | 16 % | Multi-race | 6 | 1 | 7 | <1 % |
| 45 - 49 | 441 | 20 % | Total | 1,444 | 765 | 2,209 | 100 % |
| 50 - 54 | 311 | 14 % | | | | | |
| 55 - 59 | 201 | 9 % | | | | | |
| 60+ | 159 | 7 % | | | | | |
| Total | 2,209 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|-----|-------|-------|---------|
| Men who have sex with men | 474 | 0 | 474 | 33 % |
| Injection drug use | 260 | 136 | 396 | 28 % |
| Men who have sex with men and inject drugs | 121 | 0 | 121 | 8 % |
| Heterosexual contact | 133 | 282 | 415 | 29 % |
| Transfusion/Hemophilia | 4 | 8 | 12 | 1 % |
| Perinatal | 6 | 4 | 10 | 1 % |
| Total | 998 | 430 | 1,428 | 100 % |
| <i>No Identified Risk</i> | 446 | 335 | 781 | |

Persons Newly Diagnosed with HIV/AIDS
Region 2: Baton Rouge Region

HIV CASES DIAGNOSED January - December 2008

- 301 persons were newly diagnosed with HIV in Region 2.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 175 | 58 % | Black, Non-Hispanic | 267 | 89 % |
| Women | 126 | 42 % | White, Non-Hispanic | 27 | 9 % |
| Total | 301 | 100 % | Hispanic | 4 | 1 % |
| | | | Multi-race | 3 | 1 % |
| | | | Total | 301 | 100 % |

| <u>Age at Detection</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 0 - 12 | 2 | 1 % | Men who have sex with men | 68 | 49 % |
| 13 - 24 | 72 | 24 % | Injection drug use | 14 | 10 % |
| 25 - 34 | 90 | 30 % | Men who have sex with men and inject drugs | 1 | 1 % |
| 35 - 44 | 61 | 20 % | Heterosexual contact | 53 | 38 % |
| 45+ | 76 | 25 % | Perinatal | 2 | 1 % |
| Total | 301 | 100 % | Total | 138 | 100 % |
| | | | <i>No Identified Risk</i> | <i>163</i> | |

AIDS CASES DIAGNOSED January - December 2008

- 251 persons were newly diagnosed with AIDS in Region 2.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 142 | 57 % | Black, Non-Hispanic | 214 | 85 % |
| Women | 109 | 43 % | White, Non-Hispanic | 33 | 13 % |
| Total | 251 | 100 % | Hispanic | 3 | 1 % |
| | | | Multi-race | 1 | <1 % |
| | | | Total | 251 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 13 - 24 | 29 | 12 % | Men who have sex with men | 43 | 33 % |
| 25 - 34 | 61 | 24 % | Injection drug use | 31 | 24 % |
| 35 - 44 | 68 | 27 % | Men who have sex with men and inject drugs | 6 | 5 % |
| 45+ | 93 | 37 % | Heterosexual contact | 51 | 39 % |
| Total | 251 | 100 % | Total | 131 | 100 % |
| | | | <i>No Identified Risk</i> | <i>120</i> | |

Persons Living with HIV/AIDS

Region 3: Houma Region

- **942** cumulative cases have been detected in Region 3, including **17** pediatric cases.
- **441** cumulative deaths among persons in Region 3 with HIV/AIDS have occurred, including **8** pediatric deaths.
- **635** persons are currently living with HIV/AIDS in Region 3.
Persons living with HIV: **263**
Persons living with AIDS: **372**

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|------------|--------------|------------------------|------------|------------|------------|--------------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 5 | 1 % | Black, Non-Hispanic | 214 | 145 | 359 | 57 % |
| 13 - 19 | 6 | 1 % | White, Non-Hispanic | 203 | 46 | 249 | 39 % |
| 20 - 24 | 31 | 5 % | Hispanic | 12 | 4 | 16 | 3 % |
| 25 - 29 | 61 | 10 % | Asian/Pacific Islander | 1 | 1 | 2 | <1 % |
| 30 - 34 | 69 | 11 % | American Indian | 2 | 2 | 4 | 1 % |
| 35 - 39 | 79 | 12 % | Unknown | 1 | 1 | 2 | <1 % |
| 40 - 44 | 105 | 17 % | Multi-race | 2 | 1 | 3 | <1 % |
| 45 - 49 | 116 | 18 % | Total | 435 | 200 | 635 | 100 % |
| 50 - 54 | 73 | 11 % | | | | | |
| 55 - 59 | 47 | 7 % | | | | | |
| 60+ | 43 | 7 % | | | | | |
| Total | 635 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|------------|------------|------------|--------------|
| Men who have sex with men | 217 | 0 | 217 | 52 % |
| Injection drug use | 40 | 24 | 64 | 15 % |
| Men who have sex with men and inject drugs | 13 | 0 | 13 | 3 % |
| Heterosexual contact | 32 | 81 | 113 | 27 % |
| Transfusion/Hemophilia | 0 | 1 | 1 | <1 % |
| Perinatal | 6 | 7 | 13 | 3 % |
| Total | 308 | 113 | 421 | 100 % |
| <i>No Identified Risk</i> | <i>127</i> | <i>87</i> | <i>214</i> | |

Persons Living with AIDS

Region 3: Houma Region

- **697** cumulative AIDS cases have been diagnosed in Region 3, including **9** pediatric cases.
- **372** persons are currently living with AIDS in Region 3.

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|-------|---------|------------------------|-----|-------|-------|---------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 13 - 19 | 4 | 1 % | Black, Non-Hispanic | 140 | 78 | 218 | 59 % |
| 20 - 24 | 9 | 2 % | White, Non-Hispanic | 112 | 25 | 137 | 37 % |
| 25 - 29 | 25 | 7 % | Hispanic | 8 | 2 | 10 | 3 % |
| 30 - 34 | 30 | 8 % | Asian/Pacific Islander | 0 | 1 | 1 | <1 % |
| 35 - 39 | 40 | 11 % | American Indian | 1 | 2 | 3 | 1 % |
| 40 - 44 | 68 | 18 % | Multi-race | 2 | 1 | 3 | 1 % |
| 45 - 49 | 75 | 20 % | Total | 263 | 109 | 372 | 100 % |
| 50 - 54 | 55 | 15 % | | | | | |
| 55 - 59 | 35 | 9 % | | | | | |
| 60+ | 31 | 8 % | | | | | |
| Total | 372 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|-----|-------|-------|---------|
| Men who have sex with men | 138 | 0 | 138 | 54 % |
| Injection drug use | 25 | 17 | 42 | 16 % |
| Men who have sex with men and inject drugs | 7 | 0 | 7 | 3 % |
| Heterosexual contact | 20 | 44 | 64 | 25 % |
| Transfusion/Hemophilia | 0 | 1 | 1 | <1 % |
| Perinatal | 2 | 3 | 5 | 2 % |
| Total | 192 | 65 | 257 | 100 % |
| <i>No Identified Risk</i> | 71 | 44 | 115 | |

**Persons Newly Diagnosed with HIV/AIDS
Region 3: Houma Region**

HIV CASES DIAGNOSED January - December 2008

• 46 persons were newly diagnosed with HIV in Region 3.

| <u>Gender</u> | Cases | Percent | <u>Race/Ethnicity</u> | Cases | Percent |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 28 | 61 % | Black, Non-Hispanic | 28 | 61 % |
| Women | 18 | 39 % | White, Non-Hispanic | 13 | 28 % |
| Total | 46 | 100 % | Hispanic | 3 | 7 % |
| | | | Asian/Pacific Islander | 1 | 2 % |
| | | | Unknown | 1 | 2 % |
| | | | Total | 46 | 100 % |

| <u>Age at Detection</u> | Cases | Percent | <u>Cases with a reported risk:</u> | Cases | Percent |
|-------------------------|--------------|----------------|------------------------------------|--------------|----------------|
| 13 - 24 | 10 | 22 % | Men who have sex with men | 12 | 67 % |
| 25 - 34 | 17 | 37 % | Injection drug use | 2 | 11 % |
| 35 - 44 | 11 | 24 % | Heterosexual contact | 4 | 22 % |
| 45+ | 8 | 17 % | Total | 18 | 100 % |
| Total | 46 | 100 % | <i>No Identified Risk</i> | 28 | |

AIDS CASES DIAGNOSED January - December 2008

• 33 persons were newly diagnosed with AIDS in Region 3.

| <u>Gender</u> | Cases | Percent | <u>Race/Ethnicity</u> | Cases | Percent |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 23 | 70 % | Black, Non-Hispanic | 17 | 52 % |
| Women | 10 | 30 % | White, Non-Hispanic | 13 | 39 % |
| Total | 33 | 100 % | Hispanic | 1 | 3 % |
| | | | Asian/Pacific Islander | 1 | 3 % |
| | | | Multi-race | 1 | 3 % |
| | | | Total | 33 | 100 % |

| <u>Age at Diagnosis</u> | Cases | Percent | <u>Cases with a reported risk:</u> | Cases | Percent |
|-------------------------|--------------|----------------|------------------------------------|--------------|----------------|
| 13 - 24 | 5 | 15 % | Men who have sex with men | 10 | 59 % |
| 25 - 34 | 6 | 18 % | Injection drug use | 5 | 29 % |
| 35 - 44 | 10 | 30 % | Heterosexual contact | 2 | 12 % |
| 45+ | 12 | 36 % | Total | 17 | 100 % |
| Total | 33 | 100 % | <i>No Identified Risk</i> | 16 | |

Persons Living with HIV/AIDS

Region 4: Lafayette Region

- **1,865** cumulative cases have been detected in Region 4, including **16** pediatric cases.
- **833** cumulative deaths among persons in Region 4 with HIV/AIDS have occurred, including **9** pediatric deaths.
- **1,278** persons are currently living with HIV/AIDS in Region 4.
Persons living with HIV: **585**
Persons living with AIDS: **693**

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|--------------|----------------|------------------------|------------|--------------|--------------|----------------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 3 | <1 % | Black, Non-Hispanic | 495 | 234 | 729 | 57 % |
| 13 - 19 | 12 | 1 % | White, Non-Hispanic | 395 | 84 | 479 | 37 % |
| 20 - 24 | 45 | 4 % | Hispanic | 49 | 2 | 51 | 4 % |
| 25 - 29 | 108 | 8 % | Asian/Pacific Islander | 0 | 2 | 2 | <1 % |
| 30 - 34 | 113 | 9 % | American Indian | 1 | 1 | 2 | <1 % |
| 35 - 39 | 136 | 11 % | Unknown | 2 | 0 | 2 | <1 % |
| 40 - 44 | 236 | 18 % | Multi-race | 9 | 4 | 13 | 1 % |
| 45 - 49 | 269 | 21 % | Total | 951 | 327 | 1,278 | 100 % |
| 50 - 54 | 182 | 14 % | | | | | |
| 55 - 59 | 103 | 8 % | | | | | |
| 60+ | 71 | 6 % | | | | | |
| Total | 1,278 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|------------|--------------|--------------|----------------|
| Men who have sex with men | 395 | 0 | 395 | 51 % |
| Injection drug use | 74 | 33 | 107 | 14 % |
| Men who have sex with men and inject drugs | 48 | 0 | 48 | 6 % |
| Heterosexual contact | 60 | 138 | 198 | 26 % |
| Transfusion/Hemophilia | 5 | 1 | 6 | 1 % |
| Confirmed Other | 0 | 1 | 1 | <1 % |
| Perinatal | 6 | 6 | 12 | 2 % |
| Total | 588 | 179 | 767 | 100 % |
| <i>No Identified Risk</i> | <i>363</i> | <i>148</i> | <i>511</i> | |

**Persons Living with AIDS
Region 4: Lafayette Region**

- **1,239** cumulative AIDS cases have been diagnosed in Region 4, including 9 pediatric cases.
- **693** persons are currently living with AIDS in Region 4.

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|-------|---------|------------------------|-----|-------|-------|---------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 1 | <1 % | Black, Non-Hispanic | 268 | 120 | 388 | 56 % |
| 13 - 19 | 1 | <1 % | White, Non-Hispanic | 240 | 35 | 275 | 40 % |
| 20 - 24 | 10 | 1 % | Hispanic | 22 | 0 | 22 | 3 % |
| 25 - 29 | 42 | 6 % | Asian/Pacific Islander | 0 | 1 | 1 | <1 % |
| 30 - 34 | 58 | 8 % | Multi-race | 5 | 2 | 7 | 1 % |
| 35 - 39 | 69 | 10 % | Total | 535 | 158 | 693 | 100 % |
| 40 - 44 | 137 | 20 % | | | | | |
| 45 - 49 | 158 | 23 % | | | | | |
| 50 - 54 | 111 | 16 % | | | | | |
| 55 - 59 | 64 | 9 % | | | | | |
| 60+ | 42 | 6 % | | | | | |
| Total | 693 | 100 % | | | | | |

| <u>Exposure Category</u> | | | | |
|--|-----|-------|-------|---------|
| Cases with a reported risk: | Men | Women | Total | Percent |
| Men who have sex with men | 249 | 0 | 249 | 54 % |
| Injection drug use | 52 | 17 | 69 | 15 % |
| Men who have sex with men and inject drugs | 34 | 0 | 34 | 7 % |
| Heterosexual contact | 30 | 74 | 104 | 22 % |
| Transfusion/Hemophilia | 3 | 1 | 4 | 1 % |
| Perinatal | 0 | 3 | 3 | 1 % |
| Total | 368 | 95 | 463 | 100 % |
| <i>No Identified Risk</i> | 167 | 63 | 230 | |

**Persons Newly Diagnosed with HIV/AIDS
Region 4: Lafayette Region**

HIV CASES DIAGNOSED January - December 2008

• 79 persons were newly diagnosed with HIV in Region 4.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 57 | 72 % | Black, Non-Hispanic | 51 | 65 % |
| Women | 22 | 28 % | White, Non-Hispanic | 21 | 27 % |
| Total | 79 | 100 % | Hispanic | 5 | 6 % |
| | | | Multi-race | 2 | 3 % |
| | | | Total | 79 | 100 % |

| <u>Age at Detection</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|------------------------------------|--------------|----------------|
| 13 - 24 | 14 | 18 % | Men who have sex with men | 13 | 45 % |
| 25 - 34 | 15 | 19 % | Injection drug use | 5 | 17 % |
| 35 - 44 | 16 | 20 % | Heterosexual contact | 11 | 38 % |
| 45+ | 34 | 43 % | Total | 29 | 100 % |
| Total | 79 | 100 % | <i>No Identified Risk</i> | <i>50</i> | |

AIDS CASES DIAGNOSED January - December 2008

• 64 persons were newly diagnosed with AIDS in Region 4.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 46 | 72 % | Black, Non-Hispanic | 40 | 62 % |
| Women | 18 | 28 % | White, Non-Hispanic | 22 | 34 % |
| Total | 64 | 100 % | Hispanic | 1 | 2 % |
| | | | Asian/Pacific Islander | 1 | 2 % |
| | | | Total | 64 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 13 - 24 | 3 | 5 % | Men who have sex with men | 17 | 59 % |
| 25 - 34 | 20 | 31 % | Injection drug use | 4 | 14 % |
| 35 - 44 | 20 | 31 % | Men who have sex with men and inject drugs | 1 | 3 % |
| 45+ | 21 | 33 % | Heterosexual contact | 7 | 24 % |
| Total | 64 | 100 % | Total | 29 | 100 % |
| | | | <i>No Identified Risk</i> | <i>35</i> | |

**Persons Living with HIV/AIDS
Region 5: Lake Charles Region**

- **1,192** cumulative cases have been detected in Region 5, including **9** pediatric cases.
- **442** cumulative deaths among persons in Region 5 with HIV/AIDS have occurred, including **1** pediatric death.
- **928** persons are currently living with HIV/AIDS in Region 5.
Persons living with HIV: **406**
Persons living with AIDS: **522**

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|------------|--------------|------------------------|------------|------------|------------|--------------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 3 | <1 % | Black, Non-Hispanic | 395 | 139 | 534 | 58 % |
| 13 - 19 | 16 | 2 % | White, Non-Hispanic | 236 | 60 | 296 | 32 % |
| 20 - 24 | 36 | 4 % | Hispanic | 81 | 2 | 83 | 9 % |
| 25 - 29 | 60 | 6 % | Asian/Pacific Islander | 4 | 0 | 4 | <1 % |
| 30 - 34 | 105 | 11 % | American Indian | 2 | 1 | 3 | <1 % |
| 35 - 39 | 119 | 13 % | Unknown | 1 | 0 | 1 | <1 % |
| 40 - 44 | 179 | 19 % | Multi-race | 3 | 4 | 7 | 1 % |
| 45 - 49 | 182 | 20 % | Total | 722 | 206 | 928 | 100 % |
| 50 - 54 | 131 | 14 % | | | | | |
| 55 - 59 | 50 | 5 % | | | | | |
| 60+ | 47 | 5 % | | | | | |
| Total | 928 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|------------|------------|------------|--------------|
| Men who have sex with men | 267 | 0 | 267 | 42 % |
| Injection drug use | 98 | 30 | 128 | 20 % |
| Men who have sex with men and inject drugs | 75 | 0 | 75 | 12 % |
| Heterosexual contact | 73 | 81 | 154 | 24 % |
| Transfusion/Hemophilia | 6 | 1 | 7 | 1 % |
| Perinatal | 4 | 5 | 9 | 1 % |
| Total | 523 | 117 | 640 | 100 % |
| <i>No Identified Risk</i> | <i>199</i> | <i>89</i> | <i>288</i> | |

Persons Living with AIDS
Region 5: Lake Charles Region

- **824** cumulative AIDS cases have been diagnosed in Region 5, including **5** pediatric cases.
- **522** persons are currently living with AIDS in Region 5.

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|-------|---------|------------------------|-----|-------|-------|---------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 1 | <1 % | Black, Non-Hispanic | 228 | 65 | 293 | 56 % |
| 13 - 19 | 5 | 1 % | White, Non-Hispanic | 153 | 35 | 188 | 36 % |
| 20 - 24 | 10 | 2 % | Hispanic | 36 | 0 | 36 | 7 % |
| 25 - 29 | 24 | 5 % | Asian/Pacific Islander | 2 | 0 | 2 | <1 % |
| 30 - 34 | 46 | 9 % | Multi-race | 2 | 1 | 3 | 1 % |
| 35 - 39 | 65 | 12 % | Total | 421 | 101 | 522 | 100 % |
| 40 - 44 | 118 | 23 % | | | | | |
| 45 - 49 | 113 | 22 % | | | | | |
| 50 - 54 | 78 | 15 % | | | | | |
| 55 - 59 | 33 | 6 % | | | | | |
| 60+ | 29 | 6 % | | | | | |
| Total | 522 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|-----|-------|-------|---------|
| Men who have sex with men | 173 | 0 | 173 | 44 % |
| Injection drug use | 64 | 22 | 86 | 22 % |
| Men who have sex with men and inject drugs | 50 | 0 | 50 | 13 % |
| Heterosexual contact | 35 | 42 | 77 | 19 % |
| Transfusion/Hemophilia | 5 | 1 | 6 | 2 % |
| Perinatal | 2 | 2 | 4 | 1 % |
| Total | 329 | 67 | 396 | 100 % |
| <i>No Identified Risk</i> | 92 | 34 | 126 | |

**Persons Newly Diagnosed with HIV/AIDS
Region 5: Lake Charles Region**

HIV CASES DIAGNOSED January - December 2008

• 61 persons were newly diagnosed with HIV in Region 5.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 44 | 72 % | Black, Non-Hispanic | 33 | 54 % |
| Women | 17 | 28 % | White, Non-Hispanic | 23 | 38 % |
| Total | 61 | 100 % | Hispanic | 4 | 7 % |
| | | | American Indian | 1 | 2 % |
| | | | Total | 61 | 100 % |

| <u>Age at Detection</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 13 - 24 | 14 | 23 % | Men who have sex with men | 16 | 70 % |
| 25 - 34 | 15 | 25 % | Injection drug use | 2 | 9 % |
| 35 - 44 | 17 | 28 % | Men who have sex with men and inject drugs | 1 | 4 % |
| 45+ | 15 | 25 % | Heterosexual contact | 4 | 17 % |
| Total | 61 | 100 % | Total | 23 | 100 % |
| | | | <i>No Identified Risk</i> | 38 | |

AIDS CASES DIAGNOSED January - December 2008

• 41 persons were newly diagnosed with AIDS in Region 5.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 27 | 66 % | Black, Non-Hispanic | 25 | 61 % |
| Women | 14 | 34 % | White, Non-Hispanic | 14 | 34 % |
| Total | 41 | 100 % | Hispanic | 2 | 5 % |
| | | | Total | 41 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 13 - 24 | 4 | 10 % | Men who have sex with men | 11 | 46 % |
| 25 - 34 | 14 | 34 % | Injection drug use | 3 | 12 % |
| 35 - 44 | 11 | 27 % | Men who have sex with men and inject drugs | 1 | 4 % |
| 45+ | 12 | 29 % | Heterosexual contact | 8 | 33 % |
| Total | 41 | 100 % | Perinatal | 1 | 4 % |
| | | | Total | 24 | 100 % |
| | | | <i>No Identified Risk</i> | 17 | |

**Persons Living with HIV/AIDS
Region 6: Alexandria Region**

- **1,237** cumulative cases have been detected in Region 6, including **13** pediatric cases.
- **444** cumulative deaths among persons in Region 6 with HIV/AIDS have occurred, including **4** pediatric deaths.
- **822** persons are currently living with HIV/AIDS in Region 6.
Persons living with HIV: **402**
Persons living with AIDS: **420**

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|------------|--------------|------------------------|------------|------------|------------|--------------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 2 | <1 % | Black, Non-Hispanic | 391 | 191 | 582 | 71 % |
| 13 - 19 | 4 | <1 % | White, Non-Hispanic | 145 | 47 | 192 | 23 % |
| 20 - 24 | 26 | 3 % | Hispanic | 39 | 2 | 41 | 5 % |
| 25 - 29 | 74 | 9 % | Asian/Pacific Islander | 2 | 0 | 2 | <1 % |
| 30 - 34 | 111 | 14 % | American Indian | 0 | 1 | 1 | <1 % |
| 35 - 39 | 117 | 14 % | Multi-race | 4 | 0 | 4 | <1 % |
| 40 - 44 | 134 | 16 % | Total | 581 | 241 | 822 | 100 % |
| 45 - 49 | 138 | 17 % | | | | | |
| 50 - 54 | 118 | 14 % | | | | | |
| 55 - 59 | 58 | 7 % | | | | | |
| 60+ | 40 | 5 % | | | | | |
| Total | 822 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|------------|------------|------------|--------------|
| Men who have sex with men | 195 | 0 | 195 | 37 % |
| Injection drug use | 72 | 29 | 101 | 19 % |
| Men who have sex with men and inject drugs | 54 | 0 | 54 | 10 % |
| Heterosexual contact | 63 | 99 | 162 | 31 % |
| Transfusion/Hemophilia | 9 | 1 | 10 | 2 % |
| Perinatal | 4 | 2 | 6 | 1 % |
| Total | 397 | 131 | 528 | 100 % |
| <i>No Identified Risk</i> | <i>184</i> | <i>110</i> | <i>294</i> | |

Persons Living with AIDS
Region 6: Alexandria Region

- **767** cumulative AIDS cases have been diagnosed in Region 6, including **4** pediatric cases.
- **420** persons are currently living with AIDS in Region 6.

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|-------|---------|------------------------|-----|-------|-------|---------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 13 - 19 | 1 | <1 % | Black, Non-Hispanic | 206 | 76 | 282 | 67 % |
| 20 - 24 | 5 | 1 % | White, Non-Hispanic | 94 | 28 | 122 | 29 % |
| 25 - 29 | 26 | 6 % | Hispanic | 13 | 0 | 13 | 3 % |
| 30 - 34 | 42 | 10 % | Asian/Pacific Islander | 1 | 0 | 1 | <1 % |
| 35 - 39 | 58 | 14 % | Multi-race | 2 | 0 | 2 | <1 % |
| 40 - 44 | 82 | 20 % | Total | 316 | 104 | 420 | 100 % |
| 45 - 49 | 77 | 18 % | | | | | |
| 50 - 54 | 66 | 16 % | | | | | |
| 55 - 59 | 39 | 9 % | | | | | |
| 60+ | 24 | 6 % | | | | | |
| Total | 420 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|-----|-------|-------|---------|
| Men who have sex with men | 102 | 0 | 102 | 35 % |
| Injection drug use | 37 | 17 | 54 | 19 % |
| Men who have sex with men and inject drugs | 41 | 0 | 41 | 14 % |
| Heterosexual contact | 34 | 48 | 82 | 28 % |
| Transfusion/Hemophilia | 7 | 0 | 7 | 2 % |
| Perinatal | 1 | 1 | 2 | 1 % |
| Total | 222 | 66 | 288 | 100 % |
| <i>No Identified Risk</i> | 94 | 38 | 132 | |

Persons Newly Diagnosed with HIV/AIDS

Region 6: Alexandria Region

HIV CASES DIAGNOSED January - December 2008

- 49 persons were newly diagnosed with HIV in Region 6.

| <u>Gender</u> | Cases | Percent | <u>Race/Ethnicity</u> | Cases | Percent |
|---------------|-----------|--------------|-----------------------|-----------|--------------|
| Men | 36 | 73 % | Black, Non-Hispanic | 28 | 57 % |
| Women | 13 | 27 % | White, Non-Hispanic | 11 | 22 % |
| Total | 49 | 100 % | Hispanic | 9 | 18 % |
| | | | Multi-race | 1 | 2 % |
| | | | Total | 49 | 100 % |

| <u>Age at Detection</u> | Cases | Percent | <u>Cases with a reported risk:</u> | Cases | Percent |
|-------------------------|-----------|--------------|--|-----------|--------------|
| 13 - 24 | 12 | 24 % | Men who have sex with men | 13 | 62 % |
| 25 - 34 | 17 | 35 % | Injection drug use | 1 | 5 % |
| 35 - 44 | 8 | 16 % | Men who have sex with men and inject drugs | 2 | 10 % |
| 45+ | 12 | 24 % | Heterosexual contact | 5 | 24 % |
| Total | 49 | 100 % | Total | 21 | 100 % |
| | | | <i>No Identified Risk</i> | 28 | |

AIDS CASES DIAGNOSED January - December 2008

- 36 persons were newly diagnosed with AIDS in Region 6.

| <u>Gender</u> | Cases | Percent | <u>Race/Ethnicity</u> | Cases | Percent |
|---------------|-----------|--------------|-----------------------|-----------|--------------|
| Men | 27 | 75 % | Black, Non-Hispanic | 24 | 67 % |
| Women | 9 | 25 % | White, Non-Hispanic | 8 | 22 % |
| Total | 36 | 100 % | Hispanic | 3 | 8 % |
| | | | Multi-race | 1 | 3 % |
| | | | Total | 36 | 100 % |

| <u>Age at Diagnosis</u> | Cases | Percent | <u>Cases with a reported risk:</u> | Cases | Percent |
|-------------------------|-----------|--------------|--|-----------|--------------|
| 25 - 34 | 13 | 36 % | Men who have sex with men | 6 | 29 % |
| 35 - 44 | 12 | 33 % | Injection drug use | 7 | 33 % |
| 45+ | 11 | 31 % | Men who have sex with men and inject drugs | 2 | 10 % |
| Total | 36 | 100 % | Heterosexual contact | 5 | 24 % |
| | | | Transfusion/Hemophilia | 1 | 5 % |
| | | | Total | 21 | 100 % |
| | | | <i>No Identified Risk</i> | 15 | |

**Persons Living with HIV/AIDS
Region 7: Shreveport Region**

- **1,961** cumulative cases have been detected in Region 7, including **27** pediatric cases.
- **865** cumulative deaths among persons in Region 7 with HIV/AIDS have occurred, including **10** pediatric deaths.
- **1,427** persons are currently living with HIV/AIDS in Region 7.
Persons living with HIV: **632**
Persons living with AIDS: **795**

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|--------------|--------------|------------------------|--------------|------------|--------------|--------------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 5 | <1 % | Black, Non-Hispanic | 688 | 307 | 995 | 70 % |
| 13 - 19 | 22 | 2 % | White, Non-Hispanic | 323 | 74 | 397 | 28 % |
| 20 - 24 | 89 | 6 % | Hispanic | 18 | 7 | 25 | 2 % |
| 25 - 29 | 139 | 10 % | Asian/Pacific Islander | 2 | 0 | 2 | <1 % |
| 30 - 34 | 174 | 12 % | American Indian | 4 | 0 | 4 | <1 % |
| 35 - 39 | 186 | 13 % | Unknown | 0 | 1 | 1 | <1 % |
| 40 - 44 | 226 | 16 % | Multi-race | 2 | 1 | 3 | <1 % |
| 45 - 49 | 248 | 17 % | Total | 1,037 | 390 | 1,427 | 100 % |
| 50 - 54 | 168 | 12 % | | | | | |
| 55 - 59 | 89 | 6 % | | | | | |
| 60+ | 81 | 6 % | | | | | |
| Total | 1,427 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|------------|------------|------------|--------------|
| Men who have sex with men | 498 | 0 | 498 | 51 % |
| Injection drug use | 72 | 45 | 117 | 12 % |
| Men who have sex with men and inject drugs | 111 | 0 | 111 | 11 % |
| Heterosexual contact | 81 | 150 | 231 | 24 % |
| Transfusion/Hemophilia | 4 | 3 | 7 | 1 % |
| Perinatal | 9 | 7 | 16 | 2 % |
| Total | 775 | 205 | 980 | 100 % |
| <i>No Identified Risk</i> | <i>262</i> | <i>185</i> | <i>447</i> | |

**Persons Living with AIDS
Region 7: Shreveport Region**

- **1,317** cumulative AIDS cases have been diagnosed in Region 7, including **19** pediatric cases.
- **795** persons are currently living with AIDS in Region 7.

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|-------|---------|-----------------------|-----|-------|-------|---------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 2 | <1 % | Black, Non-Hispanic | 413 | 149 | 562 | 71 % |
| 13 - 19 | 8 | 1 % | White, Non-Hispanic | 181 | 31 | 212 | 27 % |
| 20 - 24 | 25 | 3 % | Hispanic | 13 | 4 | 17 | 2 % |
| 25 - 29 | 47 | 6 % | American Indian | 2 | 0 | 2 | <1 % |
| 30 - 34 | 79 | 10 % | Multi-race | 1 | 1 | 2 | <1 % |
| 35 - 39 | 104 | 13 % | Total | 610 | 185 | 795 | 100 % |
| 40 - 44 | 138 | 17 % | | | | | |
| 45 - 49 | 161 | 20 % | | | | | |
| 50 - 54 | 118 | 15 % | | | | | |
| 55 - 59 | 66 | 8 % | | | | | |
| 60+ | 47 | 6 % | | | | | |
| Total | 795 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|-----|-------|-------|---------|
| Men who have sex with men | 284 | 0 | 284 | 49 % |
| Injection drug use | 49 | 24 | 73 | 13 % |
| Men who have sex with men and inject drugs | 69 | 0 | 69 | 12 % |
| Heterosexual contact | 60 | 82 | 142 | 24 % |
| Transfusion/Hemophilia | 3 | 3 | 6 | 1 % |
| Perinatal | 4 | 5 | 9 | 2 % |
| Total | 469 | 114 | 583 | 100 % |
| <i>No Identified Risk</i> | 141 | 71 | 212 | |

Persons Newly Diagnosed with HIV/AIDS
Region 7: Shreveport Region

HIV CASES DIAGNOSED January - December 2008

- 114 persons were newly diagnosed with HIV in Region 7.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 77 | 68 % | Black, Non-Hispanic | 93 | 82 % |
| Women | 37 | 32 % | White, Non-Hispanic | 18 | 16 % |
| Total | 114 | 100 % | Hispanic | 3 | 3 % |
| | | | Total | 114 | 100 % |

| <u>Age at Detection</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 13 - 24 | 27 | 24 % | Men who have sex with men | 33 | 62 % |
| 25 - 34 | 33 | 29 % | Injection drug use | 3 | 6 % |
| 35 - 44 | 20 | 18 % | Men who have sex with men and inject drugs | 1 | 2 % |
| 45+ | 34 | 30 % | Heterosexual contact | 16 | 30 % |
| Total | 114 | 100 % | Total | 53 | 100 % |
| | | | <i>No Identified Risk</i> | <i>61</i> | |

AIDS CASES DIAGNOSED January - December 2008

- 83 persons were newly diagnosed with AIDS in Region 7.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 60 | 72 % | Black, Non-Hispanic | 62 | 75 % |
| Women | 23 | 28 % | White, Non-Hispanic | 18 | 22 % |
| Total | 83 | 100 % | Hispanic | 2 | 2 % |
| | | | Multi-race | 1 | 1 % |
| | | | Total | 83 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 13 - 24 | 4 | 5 % | Men who have sex with men | 29 | 57 % |
| 25 - 34 | 25 | 30 % | Injection drug use | 7 | 14 % |
| 35 - 44 | 22 | 27 % | Men who have sex with men and inject drugs | 1 | 2 % |
| 45+ | 32 | 39 % | Heterosexual contact | 12 | 24 % |
| Total | 83 | 100 % | Perinatal | 2 | 4 % |
| | | | Total | 51 | 100 % |
| | | | <i>No Identified Risk</i> | <i>32</i> | |

Persons Living with HIV/AIDS
Region 8: Monroe Region

- **1,455** cumulative cases have been detected in Region 8, including **16** pediatric cases.
- **583** cumulative deaths among persons in Region 8 with HIV/AIDS have occurred, including **4** pediatric deaths.
- **950** persons are currently living with HIV/AIDS in Region 8.
Persons living with HIV: **442**
Persons living with AIDS: **508**

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|------------|--------------|------------------------|------------|------------|------------|--------------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 7 | 1 % | Black, Non-Hispanic | 439 | 291 | 730 | 77 % |
| 13 - 19 | 16 | 2 % | White, Non-Hispanic | 151 | 48 | 199 | 21 % |
| 20 - 24 | 39 | 4 % | Hispanic | 9 | 4 | 13 | 1 % |
| 25 - 29 | 102 | 11 % | Asian/Pacific Islander | 2 | 0 | 2 | <1 % |
| 30 - 34 | 129 | 14 % | American Indian | 1 | 0 | 1 | <1 % |
| 35 - 39 | 121 | 13 % | Unknown | 0 | 1 | 1 | <1 % |
| 40 - 44 | 133 | 14 % | Multi-race | 1 | 3 | 4 | <1 % |
| 45 - 49 | 151 | 16 % | Total | 603 | 347 | 950 | 100 % |
| 50 - 54 | 124 | 13 % | | | | | |
| 55 - 59 | 65 | 7 % | | | | | |
| 60+ | 63 | 7 % | | | | | |
| Total | 950 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|------------|------------|------------|--------------|
| Men who have sex with men | 273 | 0 | 273 | 44 % |
| Injection drug use | 47 | 33 | 80 | 13 % |
| Men who have sex with men and inject drugs | 40 | 0 | 40 | 6 % |
| Heterosexual contact | 62 | 148 | 210 | 34 % |
| Transfusion/Hemophilia | 3 | 3 | 6 | 1 % |
| Perinatal | 5 | 8 | 13 | 2 % |
| Total | 430 | 192 | 622 | 100 % |
| <i>No Identified Risk</i> | <i>173</i> | <i>155</i> | <i>328</i> | |

Persons Living with AIDS
Region 8: Monroe Region

- **940** cumulative AIDS cases have been diagnosed in Region 8, including **7** pediatric cases.
- **508** persons are currently living with AIDS in Region 8.

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|-------|---------|------------------------|-----|-------|-------|---------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 1 | <1 % | Black, Non-Hispanic | 253 | 127 | 380 | 75 % |
| 13 - 19 | 2 | <1 % | White, Non-Hispanic | 93 | 24 | 117 | 23 % |
| 20 - 24 | 8 | 2 % | Hispanic | 6 | 2 | 8 | 2 % |
| 25 - 29 | 32 | 6 % | Asian/Pacific Islander | 2 | 0 | 2 | <1 % |
| 30 - 34 | 68 | 13 % | Multi-race | 0 | 1 | 1 | <1 % |
| 35 - 39 | 68 | 13 % | Total | 354 | 154 | 508 | 100 % |
| 40 - 44 | 78 | 15 % | | | | | |
| 45 - 49 | 82 | 16 % | | | | | |
| 50 - 54 | 84 | 17 % | | | | | |
| 55 - 59 | 40 | 8 % | | | | | |
| 60+ | 45 | 9 % | | | | | |
| Total | 508 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|-----|-------|-------|---------|
| Men who have sex with men | 153 | 0 | 153 | 44 % |
| Injection drug use | 28 | 21 | 49 | 14 % |
| Men who have sex with men and inject drugs | 29 | 0 | 29 | 8 % |
| Heterosexual contact | 40 | 74 | 114 | 33 % |
| Transfusion/Hemophilia | 2 | 1 | 3 | 1 % |
| Perinatal | 2 | 0 | 2 | 1 % |
| Total | 254 | 96 | 350 | 100 % |
| <i>No Identified Risk</i> | 100 | 58 | 158 | |

Persons Newly Diagnosed with HIV/AIDS

Region 8: Monroe Region

HIV CASES DIAGNOSED January - December 2008

- 56 persons were newly diagnosed with HIV in Region 8.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 39 | 70 % | Black, Non-Hispanic | 46 | 82 % |
| Women | 17 | 30 % | White, Non-Hispanic | 6 | 11 % |
| Total | 56 | 100 % | Hispanic | 3 | 5 % |
| | | | Asian/Pacific Islander | 1 | 2 % |
| | | | Total | 56 | 100 % |

| <u>Age at Detection</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|------------------------------------|--------------|----------------|
| 13 - 24 | 9 | 16 % | Men who have sex with men | 25 | 74 % |
| 25 - 34 | 20 | 36 % | Heterosexual contact | 9 | 26 % |
| 35 - 44 | 12 | 21 % | Total | 34 | 100 % |
| 45+ | 15 | 27 % | <i>No Identified Risk</i> | 22 | |
| Total | 56 | 100% | | | |

AIDS CASES DIAGNOSED January - December 2008

- 43 persons were newly diagnosed with AIDS in Region 8.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 27 | 63 % | Black, Non-Hispanic | 31 | 72 % |
| Women | 16 | 37 % | White, Non-Hispanic | 9 | 21 % |
| Total | 43 | 100 % | Hispanic | 2 | 5 % |
| | | | Asian/Pacific Islander | 1 | 2 % |
| | | | Total | 43 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 13 - 24 | 2 | 5 % | Men who have sex with men | 13 | 57 % |
| 25 - 34 | 12 | 28 % | Injection drug use | 3 | 13 % |
| 35 - 44 | 16 | 37 % | Men who have sex with men and inject drugs | 1 | 4 % |
| 45+ | 13 | 30 % | Heterosexual contact | 6 | 26 % |
| Total | 43 | 100 % | Total | 23 | 100 % |
| | | | <i>No Identified Risk</i> | 20 | |

Persons Living with HIV/AIDS
Region 9: Hammond/Slidell Region

- **1,282** cumulative cases have been detected in Region 9, including **14** pediatric cases.
- **520** cumulative deaths among persons in Region 9 with HIV/AIDS have occurred, including **4** pediatric deaths.
- **962** persons are currently living with HIV/AIDS in Region 9.
 Persons living with HIV: **424**
 Persons living with AIDS: **538**

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|------------|--------------|------------------------|------------|------------|------------|--------------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 0 - 12 | 7 | 1 % | Black, Non-Hispanic | 307 | 162 | 469 | 49 % |
| 13 - 19 | 4 | <1 % | White, Non-Hispanic | 370 | 93 | 463 | 48 % |
| 20 - 24 | 39 | 4 % | Hispanic | 18 | 3 | 21 | 2 % |
| 25 - 29 | 65 | 7 % | Asian/Pacific Islander | 1 | 0 | 1 | <1 % |
| 30 - 34 | 86 | 9 % | American Indian | 1 | 0 | 1 | <1 % |
| 35 - 39 | 139 | 14 % | Unknown | 1 | 0 | 1 | <1 % |
| 40 - 44 | 167 | 17 % | Multi-race | 6 | 0 | 6 | 1 % |
| 45 - 49 | 190 | 20 % | Total | 704 | 258 | 962 | 100 % |
| 50 - 54 | 136 | 14 % | | | | | |
| 55 - 59 | 71 | 7 % | | | | | |
| 60+ | 58 | 6 % | | | | | |
| Total | 962 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|------------|------------|------------|--------------|
| Men who have sex with men | 345 | 0 | 345 | 51 % |
| Injection drug use | 67 | 49 | 116 | 17 % |
| Men who have sex with men and inject drugs | 48 | 0 | 48 | 7 % |
| Heterosexual contact | 51 | 88 | 139 | 21 % |
| Transfusion/Hemophilia | 6 | 0 | 6 | 1 % |
| Perinatal | 9 | 7 | 16 | 2 % |
| Total | 526 | 144 | 670 | 100 % |
| <i>No Identified Risk</i> | <i>178</i> | <i>114</i> | <i>292</i> | |

Persons Living with AIDS
Region 9: Hammond/Slidell Region

- **879** cumulative AIDS cases have been diagnosed in Region 9, including **6** pediatric cases.
- **538** persons are currently living with AIDS in Region 9.

| <u>Age Category</u> | | | <u>Race/Ethnicity</u> | | | | |
|---------------------|-------|---------|-----------------------|-----|-------|-------|---------|
| Current Age | Cases | Percent | | Men | Women | Total | Percent |
| 20 - 24 | 8 | 1 % | Black, Non-Hispanic | 176 | 75 | 251 | 47 % |
| 25 - 29 | 20 | 4 % | White, Non-Hispanic | 228 | 43 | 271 | 50 % |
| 30 - 34 | 37 | 7 % | Hispanic | 9 | 2 | 11 | 2 % |
| 35 - 39 | 73 | 14 % | American Indian | 1 | 0 | 1 | <1 % |
| 40 - 44 | 106 | 20 % | Multi-race | 4 | 0 | 4 | 1 % |
| 45 - 49 | 124 | 23 % | Total | 418 | 120 | 538 | 100 % |
| 50 - 54 | 85 | 16 % | | | | | |
| 55 - 59 | 45 | 8 % | | | | | |
| 60+ | 40 | 7 % | | | | | |
| Total | 538 | 100 % | | | | | |

Exposure Category

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|-----|-------|-------|---------|
| Men who have sex with men | 200 | 0 | 200 | 51 % |
| Injection drug use | 50 | 27 | 77 | 20 % |
| Men who have sex with men and inject drugs | 34 | 0 | 34 | 9 % |
| Heterosexual contact | 28 | 44 | 72 | 18 % |
| Transfusion/Hemophilia | 6 | 0 | 6 | 2 % |
| Perinatal | 2 | 1 | 3 | 1 % |
| Total | 320 | 72 | 392 | 100 % |
| <i>No Identified Risk</i> | 98 | 48 | 146 | |

Persons Newly Diagnosed with HIV/AIDS

Region 9: Hammond/Slidell Region

HIV CASES DIAGNOSED January - December 2008

- 67 persons were newly diagnosed with HIV in Region 9.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 45 | 67 % | Black, Non-Hispanic | 40 | 60 % |
| Women | 22 | 33 % | White, Non-Hispanic | 26 | 39 % |
| Total | 67 | 100 % | Unknown | 1 | 1 % |
| | | | Total | 67 | 100 % |

| <u>Age at Detection</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 0 - 12 | 1 | 1 % | Men who have sex with men | 19 | 54 % |
| 13 - 24 | 6 | 9 % | Injection drug use | 4 | 11 % |
| 25 - 34 | 21 | 31 % | Men who have sex with men and inject drugs | 1 | 3 % |
| 35 - 44 | 20 | 30 % | Heterosexual contact | 10 | 29 % |
| 45+ | 19 | 28 % | Perinatal | 1 | 3 % |
| Total | 67 | 100 % | Total | 35 | 100 % |
| | | | <i>No Identified Risk</i> | <i>32</i> | |

AIDS CASES DIAGNOSED January - December 2008

- 42 persons were newly diagnosed with AIDS in Region 9.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 32 | 76 % | Black, Non-Hispanic | 25 | 60 % |
| Women | 10 | 24 % | White, Non-Hispanic | 17 | 40 % |
| Total | 42 | 100 % | Total | 42 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk:</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|--|--------------|----------------|
| 13 - 24 | 1 | 2 % | Men who have sex with men | 13 | 57 % |
| 25 - 34 | 11 | 26 % | Injection drug use | 3 | 13 % |
| 35 - 44 | 16 | 38 % | Men who have sex with men and inject drugs | 2 | 9 % |
| 45+ | 14 | 33 % | Heterosexual contact | 4 | 17 % |
| Total | 42 | 100 % | Transfusion/Hemophilia | 1 | 4 % |
| | | | Total | 23 | 100 % |
| | | | <i>No Identified Risk</i> | <i>19</i> | |